

SG11



November 1978

*I wish we had had  
this protocol several years  
ago for our work with [unclear].*

## STANDARD REMOTE-VIEWING PROTOCOL (LOCAL TARGETS)

by  
Harold E. Puthoff and Russell Targ.

**SRI International**

STANDARD REMOTE-VIEWING PROTOCOL  
(LOCAL TARGETS)

The basic outline of our standard remote-viewing protocol is as given in our tutorial paper, "A Perceptual Channel for Information Transfer over Kilometer Distances: Historical Perspective and Recent Research," H. Puthoff and R. Targ, Proc. IEEE, pp. 329-354, March 1976.<sup>1</sup> The elements of the protocol, each of which is addressed below, consist of (1) target pool selection; (2) subject orientation; (3) outbound experimenter behavior; (4) inbound experimenter behavior; (5) post-experiment feedback; (6) judging procedure.

1. Target Pool Selection

To carry out an experimental series of, say,  $n$  trials with a subject, a list of targets  $\gg n$  should be prepared in advance by an experimenter who will not interact with the subject after that. The targets should be chosen to be distinctive, but not necessarily distinct from each other; that is, rather than just a collection of nondescript street corners one should select bridges, towers, fountains, gardens, plazas, etc., so that a judge could in principle recognize targets on the basis of correct but sketchy descriptions. On the other hand, once having chosen a fountain-type target, there should be several fountain targets; for a bridge target, several bridge targets, etc., in order to avoid the possible subject strategy of "I had a bridge yesterday, so it can't be a bridge today." The subject should be told explicitly that there are similar as well as different types of targets.

When the target list is made, each target location should be written on a card and placed in an envelope, the envelopes randomized and numbered. These should then be stored in a secure safe or similar container.

With regard to whether a target is replaced in the pool after use, the preferable procedure, from a methodological standpoint, is to replace it. (A problem with actual replacement is that the subject,

upon becoming aware of a mental image of a previous target, might be biased to reject it as memory. An acceptable alternative is to replace a used target by a new one of similar type--e.g., one fountain by another.)

## 2. Subject Orientation

Before the experiment, the subject should be shown some previous remote-viewing results with one goal in mind--to get across the idea that one should, as nearly as possible, report raw perception rather than analysis, since the former tends to be correct and the latter is almost always wrong.\* A subject needs to understand that a rounded piece of blue metal is just that, and that he should not initially try to determine what it is. Remind the subject that imagination constitutes noise in the channel, and therefore the closer he can get to raw uninterpreted imagery, the better. To have success in the above, the best guideline we have found is to choose as subjects individuals who are self-confident, uninhibited, successful, and not afraid to be wrong. No psychological test we have investigated is as reliable as the above subjective assessment in choosing subjects.

## 3. Outbound Experimenter Behavior

At the start of an experimental session, the inbound and outbound experimenters and subject should rendezvous for a relaxed informal discussion in the laboratory setting. (The outbound experimenter or experimenters must not know the target at this time.) Together they agree on a time for the subject description to start (e.g., 30 minutes hence--the length of time required to reach the furthest target in the pool; this time is then an invariant for all experiments.) The outbound experimenter then leaves the laboratory, uses a random-number generating procedure to obtain a number from 1 - n (number of targets in pool),

---

\* Figures 3 and 4 in the IEEE paper<sup>1</sup> are good examples. In Figure 4 the subject had absolutely no concept of a pedestrian overpass, but simply saw a pattern of receding squares; in Figure 3 correctly-dimensioned pools of water were misinterpreted as purification plant pools rather than recreational swimming pools.

obtains the so-numbered envelope from the target pool, and leaves the premises. (We use a Texas Instruments SR-51 hand calculator, which has a random-number function.) After driving away from the laboratory, he opens the envelope to determine the target, and then proceeds to that location. He should arrange to park and then come upon the target location at exactly the starting time so that his view of it is fresh at the beginning of the experiment. He then simply pays attention to the environment and does not let his mind wander (especially to another target). It does not appear to matter how many people comprise the outbound team, provided they do not (1) pay attention only to each other, or (2) scatter about. At the end of the agreed-upon target viewing time (usually 15 minutes) they return to the lab.

#### 4. Inbound Experimenter Behavior

During the period that the outbound experimenters spend en route to the target, the inbound experimenter and subject have a period to relax and discuss the protocols. (Inbound it is best not to have additional observers.) The goal of the inbound experimenter during this period is to make it "safe" for the subject to experience remote viewing. For the initial orientation of a new subject, this typically includes a low-key pep talk as to how remote viewing appears to be a natural, not abnormal, function, that many people appear to have done it successfully, even their first time, and always including the reminder to eschew analysis and simply render raw impressions.

Since we think that remote viewing is a difficult task, like perceiving a subliminal stimulus, we think it takes the full attentive powers of the subject. Therefore, the environment, procedures, etc., should be as natural and comfortable as possible to minimize the attention on anything other than the job at hand. No hypnosis, strobe lights, or sensory-deprivation procedures are ever used, since in our view these (novel) environmental factors take away some of the subject's much-needed attention. We are in this sense proponents of a "naturalist school." If the subject feels more comfortable smoking, or drinking a cup of coffee, that is permitted. These should be arranged ahead of time, however, so that neither subject nor experimenter leave the

experimental room while waiting for the outbound experimenter to reach his target.

The experimenter should have arranged ahead of time to have pen and paper available for drawing, and a tape recorder. When the agreed-upon experiment time arrives, the inbound experimenter simply asks the subject to "describe what impressions come to mind with regard to where the outbound experimenter is." Most subjects prefer to close their eyes, but they should simply do what comes naturally. The room lighting is preferably subdued to prevent after-image highlights, shadows on eyelids, etc. It is best that the inbound experimenter not pressure the subject to say a lot; he should act as if there is all the time in the world. Otherwise, a subject may tend to embroider descriptions just to be saying something to please the experimenter. If the subject tends toward being analytical ("I see Macy's") the experimenter must gently lead the subject into description, not analysis. ("You don't have to tell me where it is, just describe what you see.") This is the most important and difficult task of the inbound experimenter.

It is also useful for the inbound experimenter to "surprise" the subject with new viewpoints. ("Go above the scene and look down--what do you see? If you look to the left, what do you see?") The subject's viewpoint appears to shift rapidly with a question like this, and the data come through before the subject's defenses activate to block it out. The shifting of viewpoint also obviates the problem of the subject spending the entire time giving meticulous detail on a trivial item, such as a flower, which, even if true, will be of no help to a judge. Once a subject feels he sees something, he tends to hang on to this perception rather than commit himself to a new viewpoint.

The subject must be encouraged to sketch what he sees, even over his objections that he is not an artist, can't sketch, etc. He may do so throughout, or wait until the last five minutes if intermittent drawing would distract his concentration. Since drawings tend to be more accurate than verbalizations, this is an extremely important factor for good results.

5. Post-Experiment Feedback

When the outbound experimenter returns, the inbound and outbound experimenters and subject should proceed directly to the target for feedback. This helps to develop the subject's sense of which parts of his mental imaging are correct, versus incorrect. It completes the experiment for him, so that when he does a following experiment, his mind is not still involved with wondering how he did on the previous one. Only a very experienced subject can function well time after time without feedback, so this must be done for each experiment to ensure success.

6. Judging Procedure

In a sense, the most critical part of the remote-viewing procedure is the judging. Any single experiment in remote viewing, even if perfect, can in principle be dismissed as possibly coincidence. Further, any result less than perfect can be dismissed as a generalized "grass is green, sky is blue" transcript that fits every target. Only blind differential discrimination across a series of targets can put these interpretations to rest.

To prepare the transcripts for judging, an experimenter not involved in judging must read the transcripts and delete from them any reference to dates or previously used targets, so that a judge could not order the transcripts chronologically or otherwise obtain a priori information useful in matching.

Two judging procedures can then be used: Direct Matching, and Rank Ordering. Both procedures assume that n experiments have been carried out and n responses obtained. The judge must then try to determine which of the n responses goes with which of the n targets.

a. Direct-Matching Procedure

The n responses (transcripts with associated drawings) are numbered in random order and given to the judge along with the list of n targets, also in a (different) random order. The key is known by an experimenter, but not the judge. The judge then visits the target sites and constructs a one-to-one correspondence list between targets and responses without replacement; that is, no target or response is used twice.

With the correspondence list and the aid of the key, the experimenter then consults the statistical table for Direct Matching (Table 1) to determine whether the result is statistically significant. For example, if there were 5 correct matches out of 9 responses, the table indicates that the probability of obtaining such a result by chance is  $p = 0.003125$ , or roughly 3 times out of a thousand. Since the accepted standard in behavioral research is that a result can be considered significant if one obtains the value  $p \leq 0.05$ , such a result would be considered significant--that is, indicative of a nonchance correspondence.

The Direct Matching procedure is the simplest to carry out, but will give no credit for a fairly good description if a judge has difficulty in choosing between two possibilities and chooses the wrong one. This procedure is thus overly conservative. The more difficult Rank Ordering procedure, described next, gives partial credit in such a case, and is therefore a more precise statistical tool for analysis of medium-grade results.

b. Rank-Ordering Procedure

In the use of the Rank-Ordering procedure, the experimenter randomizes the targets and transcripts as before. Now, however, each of  $n$  judges is given a set of the  $n$  transcripts but only one of the target sites to investigate. Each judge's task is to visit his assigned target site, read through all the transcripts, and order them best-to-worst match (1 through 5, say, if there are five targets and five transcripts).

With the aid of the key, the experimenter then adds up the rank-ordering numbers assigned to each target's associated transcript. For example, if the actual response to a target was given a first place when a judge was looking at the target, then it gets a 1. If the actual response to a target was given a third place match when a judge was looking at that target, then it gets a 3, etc. The addition of these numbers  $1 + 3 + \dots$  then yields a number called the sum of ranks. One then consults the rank-ordering table (Table 2) for the statistic of interest. For example, if there were 5 experiments (5 targets and 5 transcripts) and the sum of ranks was 9, the table for  $5 \times 5$  gives a

Table 1

DIRECT MATCHING

THE PROBABILITIES OF M CORRECT MATCHES OUT OF N TARGETS

| M * N | 1     | 2     | 3     | 4      | 5      | 6       | 7       | 8        | 9        | 10       | 11       | 12       |
|-------|-------|-------|-------|--------|--------|---------|---------|----------|----------|----------|----------|----------|
| 0     | ...   | .5000 | .3333 | .3750  | .36667 | .36806  | .36786  | .367882  | .367879  | .367879  | .3678794 | .3678794 |
| 1     | 1.000 | ...   | .5000 | .3333  | .37500 | .36667  | .36806  | .367857  | .367882  | .367879  | .3678795 | .3678794 |
| 2     | .5000 | ...   | .2500 | .2500  | .16667 | .18750  | .18333  | .184028  | .183929  | .183941  | .1839396 | .1839397 |
| 3     | .1667 | ...   | .1667 | ...    | .08333 | .05556  | .06250  | .061111  | .061343  | .061310  | .0613137 | .0613132 |
| 4     |       |       |       | .0417  | ...    | .02083  | .01389  | .015625  | .015278  | .015336  | .0153274 | .0153284 |
| 5     |       |       |       | .00833 | ...    | .00417  | .002778 | .003125  | .003056  | .0030671 | .0030671 | .0030655 |
| 6     |       |       |       | .00139 | ...    | .000694 | .000463 | .000521  | .0005093 | .0005112 | .0005112 | .0005112 |
| 7     |       |       |       |        | .00020 | ...     | .000099 | .000066  | .0000744 | .0000728 | .0000728 | .0000728 |
| 8     |       |       |       |        |        | .000025 | ...     | .000012  | .0000083 | .0000093 | .0000093 | .0000093 |
| 9     |       |       |       |        |        | .000003 | ...     | .0000014 | .0000009 | .0000009 | .0000009 | .0000009 |
| 10    |       |       |       |        |        |         | .000000 | ...      | .0000001 | .0000001 | .0000001 | .0000001 |
| 11    |       |       |       |        |        |         |         | .0000000 | ...      | .0000000 | .0000000 | .0000000 |
| 12    |       |       |       |        |        |         |         |          |          |          |          | .0000000 |

SIGNIFICANT AT  $p \leq 0.05$   
(4 or more out of any arbitrary N)



Table 2

## RANK-ORDERING TABLE

Number of Targets = 4; Number of Transcripts = 4

| SUM OF RANKS | P-VALUE*    |
|--------------|-------------|
| 4            | 0.39063E-02 |
| 5            | 0.19531E-01 |
| 6            | 0.58594E-01 |
| 7            | 0.13672E 00 |
| 8            | 0.25781E 00 |
| 9            | 0.41406E 00 |
| 10           | 0.58594E 00 |
| 11           | 0.74219E 00 |
| 12           | 0.86328E 00 |
| 13           | 0.94141E 00 |
| 14           | 0.98047E 00 |
| 15           | 0.99609E 00 |
| 16           | 0.10000E 01 |

Number of Targets = 5; Number of Transcripts = 5

| SUM OF RANKS | P-VALUE     |
|--------------|-------------|
| 5            | 0.32000E-03 |
| 6            | 0.19200E-02 |
| 7            | 0.67200E-02 |
| 8            | 0.17920E-01 |
| 9            | 0.40320E-01 |
| 10           | 0.79040E-01 |
| 11           | 0.13824E 00 |
| 12           | 0.21984E 00 |
| 13           | 0.32224E 00 |
| 14           | 0.43904E 00 |
| 15           | 0.56096E 00 |
| 16           | 0.67776E 00 |
| 17           | 0.78016E 00 |
| 18           | 0.86176E 00 |
| 19           | 0.92096E 00 |
| 20           | 0.95968E 00 |
| 21           | 0.98208E 00 |
| 22           | 0.99328E 00 |
| 23           | 0.99808E 00 |
| 24           | 0.99968E 00 |
| 25           | 0.10000E 01 |

---

\*The notation E-02 is to be understood as  $10^{-2}$ ; E 01 as  $10^1$ ; etc.

Table 2 (Continued)

Number of Targets = 6; Number of Transcripts = 6

| SUM OF RANKS | P-VALUE     |
|--------------|-------------|
| 6            | 0.21433E-04 |
| 7            | 0.15003E-03 |
| 8            | 0.60014E-03 |
| 9            | 0.18004E-02 |
| 10           | 0.45010E-02 |
| 11           | 0.99023E-02 |
| 12           | 0.19676E-01 |
| 13           | 0.35880E-01 |
| 14           | 0.60764E-01 |
| 15           | 0.96472E-01 |
| 16           | 0.14463E 00 |
| 17           | 0.20585E 00 |
| 18           | 0.27939E 00 |
| 19           | 0.36310E 00 |
| 20           | 0.45357E 00 |
| 21           | 0.54642E 00 |
| 22           | 0.63689E 00 |
| 23           | 0.72061E 00 |
| 24           | 0.79415E 00 |
| 25           | 0.85537E 00 |
| 26           | 0.90353E 00 |
| 27           | 0.93923E 00 |
| 28           | 0.96412E 00 |
| 29           | 0.98032E 00 |
| 30           | 0.99010E 00 |
| 31           | 0.99550E 00 |
| 32           | 0.99820E 00 |
| 33           | 0.99940E 00 |
| 34           | 0.99985E 00 |
| 35           | 0.99998E 00 |
| 36           | 0.10000E 01 |

Table 2 (Continued)

Number of Targets = 7; Number of Transcripts = 7

| SUM OF RANKS | P-VALUE     |
|--------------|-------------|
| 7            | 0.12143E-05 |
| 8            | 0.97141E-05 |
| 9            | 0.43714E-04 |
| 10           | 0.14571E-03 |
| 11           | 0.40071E-03 |
| 12           | 0.96170E-03 |
| 13           | 0.20837E-02 |
| 14           | 0.41589E-02 |
| 15           | 0.77458E-02 |
| 16           | 0.13585E-01 |
| 17           | 0.22595E-01 |
| 18           | 0.35838E-01 |
| 19           | 0.54453E-01 |
| 20           | 0.79544E-01 |
| 21           | 0.11205E 00 |
| 22           | 0.15259E 00 |
| 23           | 0.20137E 00 |
| 24           | 0.25802E 00 |
| 25           | 0.32161E 00 |
| 26           | 0.39065E 00 |
| 27           | 0.46315E 00 |
| 28           | 0.53685E 00 |
| 29           | 0.60935E 00 |
| 30           | 0.67839E 00 |
| 31           | 0.74198E 00 |
| 32           | 0.79863E 00 |
| 33           | 0.84741E 00 |
| 34           | 0.88795E 00 |
| 35           | 0.92045E 00 |
| 36           | 0.94555E 00 |
| 37           | 0.96416E 00 |
| 38           | 0.97740E 00 |
| 39           | 0.98641E 00 |
| 40           | 0.99225E 00 |
| 41           | 0.99584E 00 |
| 42           | 0.99791E 00 |
| 43           | 0.99903E 00 |
| 44           | 0.99958E 00 |
| 45           | 0.99984E 00 |
| 46           | 0.99995E 00 |
| 47           | 0.99998E 00 |
| 48           | 0.99999E 00 |
| 49           | 0.10000E 01 |

Table 2 (Continued)

Number of Targets = 8; Number of Transcripts = 8

| SUM OF RANKS | P-VALUE     |
|--------------|-------------|
| 8            | 0.59605E-07 |
| 9            | 0.53644E-06 |
| 10           | 0.26822E-05 |
| 11           | 0.98348E-05 |
| 12           | 0.29504E-04 |
| 13           | 0.76711E-04 |
| 14           | 0.17899E-03 |
| 15           | 0.38356E-03 |
| 16           | 0.76663E-03 |
| 17           | 0.14447E-02 |
| 18           | 0.25867E-02 |
| 19           | 0.44264E-02 |
| 20           | 0.72724E-02 |
| 21           | 0.11515E-01 |
| 22           | 0.17628E-01 |
| 23           | 0.26157E-01 |
| 24           | 0.37702E-01 |
| 25           | 0.52890E-01 |
| 26           | 0.72328E-01 |
| 27           | 0.96562E-01 |
| 28           | 0.12602E 00 |
| 29           | 0.16095E 00 |
| 30           | 0.20139E 00 |
| 31           | 0.24714E 00 |
| 32           | 0.29772E 00 |
| 33           | 0.35237E 00 |
| 34           | 0.41012E 00 |
| 35           | 0.46982E 00 |
| 36           | 0.53018E 00 |
| 37           | 0.58988E 00 |
| 38           | 0.64763E 00 |
| 39           | 0.70228E 00 |
| 40           | 0.75286E 00 |
| 41           | 0.79860E 00 |
| 42           | 0.83905E 00 |
| 43           | 0.87398E 00 |
| 44           | 0.90344E 00 |
| 45           | 0.92767E 00 |
| 46           | 0.94711E 00 |
| 47           | 0.96229E 00 |
| 48           | 0.97384E 00 |
| 49           | 0.98237E 00 |
| 50           | 0.98849E 00 |
| 51           | 0.99273E 00 |
| 52           | 0.99557E 00 |
| 53           | 0.99741E 00 |

Table 2 (Continued)

Number of Targets = 9; Number of Transcripts = 9

| SUM OF RANKS | P-VALUE     |
|--------------|-------------|
| 9            | 0.25812E-08 |
| 10           | 0.25812E-07 |
| 11           | 0.14196E-06 |
| 12           | 0.56786E-06 |
| 13           | 0.18455E-05 |
| 14           | 0.51675E-05 |
| 15           | 0.12919E-04 |
| 16           | 0.29529E-04 |
| 17           | 0.62748E-04 |
| 18           | 0.12547E-03 |
| 19           | 0.23821E-03 |
| 20           | 0.43226E-03 |
| 21           | 0.75357E-03 |
| 22           | 0.12673E-02 |
| 23           | 0.20628E-02 |
| 24           | 0.32586E-02 |
| 25           | 0.50075E-02 |
| 26           | 0.75003E-02 |
| 27           | 0.10968E-01 |
| 28           | 0.15683E-01 |
| 29           | 0.21954E-01 |
| 30           | 0.30122E-01 |
| 31           | 0.40548E-01 |
| 32           | 0.53601E-01 |
| 33           | 0.69639E-01 |
| 34           | 0.88989E-01 |
| 35           | 0.11192E 00 |
| 36           | 0.13864E 00 |
| 37           | 0.16924E 00 |
| 38           | 0.20370E 00 |
| 39           | 0.24189E 00 |
| 40           | 0.28353E 00 |
| 41           | 0.32821E 00 |
| 42           | 0.37540E 00 |
| 43           | 0.42447E 00 |
| 44           | 0.47469E 00 |
| 45           | 0.52531E 00 |
| 46           | 0.57553E 00 |
| 47           | 0.62460E 00 |
| 48           | 0.67179E 00 |
| 49           | 0.71647E 00 |
| 50           | 0.75811E 00 |
| 51           | 0.79630E 00 |
| 52           | 0.83076E 00 |
| 53           | 0.86136E 00 |
| 54           | 0.88807E 00 |

Table 2 (Concluded)

Number of Targets = 10; Number of Transcripts = 10

| SUM OF RANKS | P-VALUE     |
|--------------|-------------|
| 10           | 0.10000E-09 |
| 11           | 0.11000E-08 |
| 12           | 0.66000E-08 |
| 13           | 0.28600E-07 |
| 14           | 0.10010E-06 |
| 15           | 0.30030E-06 |
| 16           | 0.80080E-06 |
| 17           | 0.19448E-05 |
| 18           | 0.43758E-05 |
| 19           | 0.92378E-05 |
| 20           | 0.18475E-04 |
| 21           | 0.35261E-04 |
| 22           | 0.64599E-04 |
| 23           | 0.11412E-03 |
| 24           | 0.19512E-03 |
| 25           | 0.32387E-03 |
| 26           | 0.52317E-03 |
| 27           | 0.82418E-03 |
| 28           | 0.12686E-02 |
| 29           | 0.19106E-02 |
| 30           | 0.28197E-02 |
| 31           | 0.40825E-02 |
| 32           | 0.58049E-02 |
| 33           | 0.81133E-02 |
| 34           | 0.11156E-01 |
| 35           | 0.15103E-01 |
| 36           | 0.20143E-01 |
| 37           | 0.26484E-01 |
| 38           | 0.34347E-01 |
| 39           | 0.43960E-01 |
| 40           | 0.55552E-01 |
| 41           | 0.69345E-01 |
| 42           | 0.85541E-01 |
| 43           | 0.10432E 00 |
| 44           | 0.12581E 00 |
| 45           | 0.15011E 00 |
| 46           | 0.17725E 00 |
| 47           | 0.20721E 00 |
| 48           | 0.23987E 00 |
| 49           | 0.27506E 00 |
| 50           | 0.31255E 00 |
| 51           | 0.35202E 00 |
| 52           | 0.39311E 00 |
| 53           | 0.43538E 00 |
| 54           | 0.47838E 00 |

probability of obtaining such a rank ordering result by chance of 0.0403..., which is significant. A more complete set of tables is given in Solfvin et al.<sup>2</sup>

REFERENCES

1. H.E. Puthoff and R. Targ, "A Perceptual Channel for Information Transfer over Kilometer Distances: Historical Perspective and Recent Research," Proc. IEEE, Vol. 64, pp. 329-354 (March 1976).
2. G. Solfvin et al., "Some New Methods of Analysis for Preferential-Ranking Data," J. Am. Soc. for Psychical Research, Vol. 72, No. 2 (April 1978).